



DASKA BARNETT
PRINCIPAL

BSc. (Hons) MCOptom

ELAINE STYLES

BSc. (Hons) MCOptom Dip CLP

CHLOE ROBSON

BSc. (Hons) MCOptom

OPTOMETRISTS – CONTACT LENSES

OCULAR HYPERTENSION

What is ocular hypertension?

Ocular hypertension is when the intraocular pressure is higher than normal in the absence of optic nerve damage or visual field loss. Normal intraocular pressure (IOP) is between 10mmHg and 21mmHg. Raised IOP is one of the risk factors of Glaucoma, so those with ocular hypertension are considered to be at a higher risk of developing Glaucoma, which is why it is most important for people with **ocular hypertension** to be monitored carefully in order that any glaucoma that does develop, is detected at the earliest possible stage when treatment is most effective.

The importance of corneal thickness

Recent discoveries about the cornea, the clear part of the eye's protective covering, are showing that corneal thickness is an important factor in accurately diagnosing eye pressure. When we measure the IOP we don't always get a truly accurate reading. The IOP reading depends on the thickness of the cornea, thicker corneas cause falsely higher eye pressure readings and thinner corneas cause falsely lower eye pressure readings. It is also known that people with a thin central cornea less than 0.555mm are significantly more at risk of developing Primary Open Angle Glaucoma.

Your corneal thickness therefore affects your risk for glaucoma, and knowing what your corneal thickness is can make your diagnosis more accurate. With an ultrasonic device called a Pachymeter we can measure your corneal thickness, from this we can calculate your true IOP.

What is Glaucoma?

Glaucoma is the name for a group of eye conditions in which the optic nerve is damaged at the point where it leaves the eye. This causes a loss of peripheral vision which is why it can go undetected until the later stages. If it remains undetected for a long period of time then it may result in "tunnel vision" and blindness.

In the UK some form of glaucoma affects about 2 in 100 people over the age of 40. If detected early enough, glaucoma can usually be treated. In most cases, eyedrops to reduce the pressure will be prescribed, although in some cases a minor operation is needed. Treatment is unable to restore vision that has been lost but aims to preserve the remaining sight.

Regular eye examinations where the ocular health, eye pressure and field of vision are checked are essential. Raised IOP is NOT the same as high blood pressure. If you are worried about glaucoma, or would like any more information just contact the practise and we will be happy to talk to you.

36 EARLS COURT ROAD LONDON W8 6EA TEL: 020 7938 1251 FAX: 7938 4403

EMAIL: info@specsofken.com WEBSITE: www.specsofken.com